



Forticlina retard

Long-Acting Oxytetracycline



Pharmaceutical Form

Injectable solution.

Composition per ml

Oxytetracycline

200 mg

Indications

Treatment of germs sensitive to oxytetracycline, which cause infections such as:

Cattle: pneumonia, shipping fever, pododermatitis, bacterial enteritis, actinobacillosis, omphalitis, metritis, mastitis, infected wounds and anaplasmosis.

Pigs: pneumonia, bacterial enteritis, haemorrhagic septicaemia, erysipelas and MMA syndrome.

Sheep and goats: pneumonia, chlamydia infections, metritis, mastitis, foot rot, omphalitis and infected wounds.

Administration Route

Deep intramuscular injection.

Dosage and administration

Cattle: 1 ml per 10 kg body weight. Do not apply more than 15 ml in the same injection site.

Pigs: 1 ml per 10 kg body weight by deep intramuscular injection. Do not apply more than 10 ml per injection site. In piglets smaller than 10 kg administer 1 ml per animal subcutaneously.

Sheep, Goats: 1 ml per 10 kg body weight. Do not apply more than 5 ml per injection site.

At the indicated dosage the therapeutic level stays in blood for 3-4 days. Therefore there is no need to repeat the treatment unless the veterinarian ask to do so.

Withdrawal period

Meat: Cattle: 30 days, Swine: 23 days, Sheep and goats: 24 days.

Milk: Do not use in dairy animals whose milk is for human consumption.

Shelf-life period

3 years.

Storage

Store in a cool, dark place.

Supply

Vials of 100 ml and 250 ml.